

Forage Maize Varieties



Irish Recommended List 2016

CROPS EVALUATION AND CERTIFICATION DIVISION

Forage Maize Varieties

Irish Recommended List 2016

CROPS EVALUATION & CERTIFICATION DIVISION

CONTENTS

	Page
Maize Growing in Ireland.	2
Introduction.	2
Variety Testing Procedure.	3
Laboratory Analysis.	4
Explanation of Results.	4
Recommended List (2016):	
Table 1: Varieties suitable for growing with plastic cover.	6
Characteristics of varieties suitable for growing with plastic cover.	7
Appendices:	
Appendix 1: Varieties evaluated in plastic covered maize National List/ Recommended List 2015.	9
Appendix 2: 2015 Recommended List of varieties suitable for growing without plastic cover and associated characteristics.	10

IMPORTANT NOTICE

The Department of Agriculture, Food and the Marine (DAFM) has taken all due care in evaluating the performance of the listed varieties for yield, quality, disease resistance and the important agronomic characters over a wide range of soils and environmental conditions, for a minimum period of 3 years. The Department cannot, however, accept responsibility for any loss or inconvenience arising from any future variation in absolute or relative varietal performance.

ACKNOWLEDGEMENTS

The Department of Agriculture, Food and the Marine wishes to acknowledge the help and co-operation received from Maize breeders, Seed Agents, Agricultural advisors with Teagasc and the growers on whose land the trials were conducted.

Maize Growing in Ireland

Forage maize grows best in areas of Ireland with a long growing season, freedom from late spring frosts and early winter frosts, high overall temperatures and maximum sunlight. Fields chosen for maize production should be south facing where possible in order to maximise sunlight and temperatures over the growing season. Elevated and exposed sites are generally unsuitable. Maize needs good soil and is not tolerant of soil compaction, so care must be taken to avoid this problem. Very light sandy soils will give reduced yields. Growing on very heavy clay soils, particularly if they have underlying poor drainage, can delay sowing date and make harvesting difficult as well as causing damage to the soil structure. In Ireland, maize is sown in spring and is grown under two production systems; 1. *'Without plastic cover' (Uncovered)*, and 2. *'With Plastic Cover'*. In recent years there has been a large swing to growing Maize under plastic with an estimated 80% of the crop currently grown in this system. Larger variability in yields and quality of uncovered crops compared to relatively consistent yields and quality in plastic covered crops from year to year, which reduce the adverse effects of poor climatic conditions has largely contributed to this move towards plastic covered crops.

The plastic, in the form of a thin sheet spread over two crop rows, is applied to the soil surface at the time of sowing. The plastic cover allows sowing to be carried out earlier than for uncovered crops, because the plastic covering quickly brings about a significant increase in the temperature of the air and soil beneath it.

Sowing dates should be decided on the basis of the usual climate and the experience gained in the area over the years.

The 2016 Maize Recommended List is based on data collected from DAFM trials from 2013 to 2015. The average sowing and harvesting dates for these trials was 16th April and 19th October respectively.

Introduction

This booklet lists the forage maize varieties that are considered most suitable for growing under Irish conditions. Reflecting the move towards growing maize under plastic cover, DAFM discontinued Recommended List (RL) trialling of uncovered Maize in 2015 and for that reason this publication does not include an updated uncovered RL. The last uncovered Maize RL from 2015 is included in the appendix to this booklet.

The varieties included on the Recommended List have completed a minimum of three years in the Department of Agriculture, Food and the Marine trials on farms situated in various geographic locations considered suitable for maize production. The trials are grown within commercial maize crops and in accordance with good farming practice.

The trials were located in Cork, South Kilkenny, Meath and Kildare during the period 2013 to 2015. The yearly trial results over these three years are combined in a single evaluation. All varieties trialled in 2015 are listed in the appendix to this booklet.

Variety Testing Procedure

New varieties are submitted annually to the Department from Irish Agents acting on behalf of International plant breeders or directly from these breeders. These varieties enter combined National List/ Recommended List trials. Trials comprise 18 varieties each year and are grown at four locations. The varieties are assessed as to their suitability under Irish conditions for dry matter yield, dry matter content, starch content and other traits.

After completing 3 years in official trials, new varieties showing superior performance are given positive Value for Cultivation and Use (V.C.U.) status. Those varieties can be National Listed by the breeder provided they have been awarded a DUS certificate, thus allowing them to be considered for Provisional Recommendation. If these Provisionally Recommended varieties continue to perform well, they may be upgraded to Full Recommendation status after 1, 2 or occasionally 3 years of further trialling. Provisional recommendation is possible for a maximum of three years.

Growers should give preference to the varieties listed unless there is compelling evidence that other varieties are more suited to their specific conditions or requirements.

Throughout the trial programme, efforts are being made to select better and earlier maturing varieties that will improve the yield and quality of the crop, as well as allowing it to be successfully grown in areas that were previously considered climatically marginal for forage maize production.

Types of Recommendation

Varieties appearing on the list for the first time are **Provisionally Recommended (PR)**, and have completed three years in combined National list/ Recommended list (NL/RL) trials. The eventual status of these varieties is determined by their subsequent level of performance in ongoing NL/RL trials; fully **Recommended (R)** classification may be deemed to be merited after a further one, two or exceptionally three years, or alternatively they may be removed from the list at any stage.

Laboratory Analysis

The Department of Agriculture, Food and the Marine take whole-crop samples from each trial plot at harvesting and dry them to determine the dry matter content. These dried samples are analysed by FBA Laboratories Ltd., Cappoquin, Co. Waterford and SAC Laboratories in Scotland..

Explanation of Results

Yield of Dry Matter/ha: Dry matter yield per hectare is calculated by measuring the fresh yield of the crop, eliminating the moisture content and expressing results on a per hectare basis. Table 1 shows the average dry matter yield of the control varieties in the 2016 DAFM Recommend List trials was 18.7t/ha. In the attached table relative yields are used to show differences between varieties with the average of the control varieties set at 100. Varieties with a score in excess of 100 achieved higher yields and varieties with a score of less than 100 achieved lower yields than the controls.

Dry matter yield of forage maize is important as this is the portion of the crop which contains the nutrient content. In assessing forage maize varieties for yield, dry matter yields should always be used instead of fresh yields as differences in the dry matter/moisture of the crop may result in fresh yields that are not correlated in any way with dry matter yield.

Starch Content: In the lead up to maturity, sugars in the plant move to the cob where they are converted to starch. Starch is an important indicator of the feeding value of the crop and is the main source of energy content in a well matured crop at harvest. Starch content is expressed as a percentage of the dry matter and is determined by wet chemistry. Table 1 shows the average starch content of the control varieties in the 2016 DAFM Recommend List trials was 26.3%. Relative yields are used to show differences between varieties with the average of the control varieties set at 100. Varieties with a score in excess of 100 had higher starch content and varieties with a score of less than 100 had lower starch content than the controls.

ME (MJ/kg): The ME system is used for differentiating maize varieties instead of the net energy system now commonly in use because the net energy system does not currently allow us to compare different maize varieties on their nutritional value. The ME value is an estimate of the energy available to an animal as a result of digestion of the feed material and is expressed in Megajoules per Kilogram.

Plant Height: Plant Height is measured from ground level to the point where the uppermost leaf departs the plant stem

Year first Recommended: This informs us of the first year the variety has first been recommended.

Dry Matter Content/Earliness of Maturity: A high dry matter is a good indicator of how much the crop has matured prior to harvest. The figures presented in Table 1 are on a relative basis, with the average of the control varieties set at 100, figures in excess of 100 have a higher dry matter content and figures less than 100 have a lower dry matter content.

For the 2016 Recommended List, the relative dry matter figure has an associated text, which indicates the maturity of the variety (Extremely Early, Very Early, Early, Medium-Early, Medium-Late, Late and Very Late). This text is used as an aid to growers in the selection of maize varieties suitable for their circumstances. Maize is a marginal crop for growing in Ireland with large differences in the favourability of maize growing sites. Therefore selection of varieties that are suited to specific sites is an important decision when growing maize in Ireland.

A maize grower in a less favourable site would generally be recommended to select an earlier maturing variety particularly if sowing late in the season. A maize grower in a more favourable site would generally be recommended to select a late maturing variety particularly if sowing early in the season. Late maturing varieties are generally higher yielding than early maturing varieties. However, starch content is an important consideration which must also be taken into account and late maturing varieties should not be sown in less favourable sites as this would result in crops not maturing and having a low starch content, particularly in poor maize growing years.

Table 1: Recommended List (2016) of Forage Maize varieties suitable for growing with plastic cover (Covered)

Actual yield data is shown for the mean of the control varieties, and the relative yield data (as % of controls) is shown for all varieties. The data is based on results of trials carried out in the period 2013 to 2015.

	Yield of Dry Matter (t/ha)	Starch content (%)	ME (MJ/kg)	Plant Height (metres)	Year first Recommended	Dry Matter Content / Earliness of Maturity
Controls*	18.7	26.3	—	—	—	35.2
Ambition (PR-1)	94	120	11.6	2.07	2016	119 Extremely Early
Award (R)	102	104	11.4	2.10	2011	102 Medium-Early
Galbi CS (PR-1)	108	90	10.9	2.25	2016	92 Late
Grosso (PR-2)	98	104	11.3	2.23	2015	93 Medium-Late
Justina (R)	100	97	11.3	2.23	2005	97 Medium-Late
LG30211 (PR-1)	101	107	11.4	2.16	2016	98 Medium-Early
P7905 (R)	105	100	11.5	2.16	2015	95 Medium-Late
Tekni (R)	98	100	11.3	2.12	2010	101 Medium-Early

* The Control varieties were Award, Justina & Tekni in 2013, 2014 and 2015.

(R): Recommended for general use. (PR): Provisionally Recommended (The number after the PR indicates the number of years provisionally recommended).

Characteristics of the varieties in Table 1 when grown with plastic cover.

- Ambition (PR-1): With Plastic:** First entered onto the plastic covered list in 2016 with a provisional recommendation. This is the lowest yielding variety on the list. It has the highest starch content on the list. It has extremely high dry matter content and is an extremely early maturing variety. It is suitable for late sowing and/or early harvest particularly on less favourable sites. Bred by Limagrain Verneuil Holding, France. Irl Agent: Goldcrop Ltd
- Award (R): With Plastic:** A good yielding variety. Starch content is very high. Dry matter content is good and it is a medium-early maturing variety. Bred by Limagrain Advanta Nederland BV, The Netherlands. Irl Agent: Goldcrop Ltd.
- Galbi CS (PR-1): With Plastic:** First entered onto the plastic covered list in 2016 with a provisional recommendation. This is the highest yielding variety on the list. It has the lowest starch content on the list. Dry matter content is low and it is the latest maturing variety on the list. It should only be sown early on favourable sites. Bred by Caussade Semences, France. Irl Agent: Goldcrop Ltd.
- Grosso (PR-2): With Plastic:** First entered the list in 2015 with a provisional recommendation. A good yielding variety with high starch content. It has moderate dry matter content and is a medium-late maturing variety. More suited for early sowing in favourable sites than in unfavourable sites. Bred by KWS Saat, Germany. Irl Agent: DLF Seeds.
- Justina (R): With Plastic:** A good yielding variety with moderate starch content. Dry matter content is moderate and it is a medium-late maturing variety. Bred by DuPont Pioneer, Germany.
- LG30211 (PR-1): With Plastic:** First entered onto the list in 2016 with a provisional recommendation. A good yielding variety with high starch content. Dry matter content is good and it is a medium-early maturing variety. Bred by Limagrain Verneuil Holding, France. Irl Agent: DLF Seeds

P7905 (R): **With Plastic:** First entered the list in 2015 with a provisional recommendation and has been upgraded to a fully recommended variety in 2016. A high yielding variety with good starch content. It has moderate dry matter content and is a medium-late maturing variety. More suited to early sowing in favourable sites than in unfavourable sites. Bred by DuPont Pioneer, Germany.

Tekni (R): **With Plastic:** A good yielding variety with good starch content. It has good dry matter content and is a medium-early maturing variety. Bred by Caussade Semences, France. Irl Agent: Goldcrop Ltd.

Appendix 1: Varieties evaluated in the Plastic Covered Maize National List/ Recommended List Trial 2015

	Variey / Breeder's Reference	Trial Year '15	Agent	Breeder & Country	Breeder's Reference
1	Ambition	3	GC	Lim. Vern (F)	LZM159/87
2	Award	8	GC	Lim. Vern. (F)	ADV 4635
3	Corfinio	2	DLF	KWS Saat (DE)	KXB 3151
4	Galbi CS	3	GC	Caussade (F)	CSM 6107
5	Grosso	4	DLF	KWS Saat (DE)	KXA 8151
6	Justina	14	DP	DuPont Pioneer	X0806A
7	KXB 3181	1	DLF	KWS Saat (DE)	KXB 3181
8	KXB 4222	1	DLF	KWS Saat (DE)	KXB 4222
9	LG30211	3	DLF	Lim. Vern (F)	LZM158/72
10	LG30.212	2	GC	Lim. Vern (F)	LZM162/33
11	MAS 20S	1	DLF	Mais. Semc (F)	MGM231407
12	P7905	4	DP	DuPont Pioneer	X8V245
13	P8200	2	DP	DuPont Pioneer	X75B142
14	P8201	1	DP	DuPont Pioneer	X80D062
15	Schobbi CS	3	GC	Caussade (F)	CSM 0163
16	Spyci CS	1	GC	Caussade (F)	CSM 2152
17	SY Feeditop	2	GC	Syngenta (F)	SB 0850
18	Tekni CS	9	GC	Caussade (F)	CSM 3122

Varieties in **bold text** are listed on the 2016 Plastic Covered Recommended List

Trial Year '15 indicates number of years the variety has been in trial

Agent: GC = Goldcrop, DLF = DLF Seeds, DP = DuPont Pioneer

Breeder: 'Lim. Vern.' indicates 'Limagrain Verneuil Holding', 'Caussade' indicates 'Caussade Semences', 'KWS Saat' indicates 'KWS Saat AG', 'Mais. Semc' indicates 'Maisadour Semences' and 'Satb Linz' indicates 'Saatbau Linz GmbH',

Appendix 2

Recommended List (2015) of Forage Maize varieties suitable for growing without plastic cover (Uncovered)

Actual yield data is shown for the mean of the control varieties, and the relative yield data (as % of controls) is shown for all varieties. The data is based on results of trials carried out over three years in the period 2012 to 2014.

	Yield of Dry Matter (t/ha)	Dry Matter content (%)	Starch content (%)	ME (MJ/kg)	Plant Height (metres)	Year first Recommended
Controls*	15.5t/ha	35.4	23.3	—	—	—
Activate (PR-1)	96	120	117	11.31	1.96	2015
Ambition (R)	105	106	113	11.33	2.15	2014
Atrium (R)	101	90	105	11.12	2.01	2013
Beacon (R)	99	99	108	11.08	2.13	2012
Beethoven (R)	104	94	88	10.96	2.15	2010
Kroft (R)	92	113	121	11.08	1.97	2014
Severus (PR-1)	102	104	114	10.90	2.06	2015

The Control varieties were Andante, Fergus and Destiny in 2012 trials, Beethoven, Beacon and Destiny in 2013 trials and Beethoven and Beacon in 2014 trials.

* Due to insufficient lodging occurring during the trial period 2012 to 2014, it was not possible to provide varietal lodging data.

(R): Recommended for general use. (PR): Provisionally Recommended (The number after the PR, indicates the number of years provisionally recommended).

Yield and quality data shown for Uncovered trials and Plastic covered trials are not directly comparable, because the trials were grown in different locations and under different conditions.

Characteristics of the varieties in Appendix 2 when grown without plastic cover

- Ambition (R):** **Uncovered:** Fully Recommended variety in 2015. High relative yield and dry matter with very high starch content. An early maturing variety. Bred by Limagrain Verneuil Holding, France. Irl Agent: Goldcrop Ltd.
- Atrium (R):** **Uncovered:** Good yielding variety. It has the lowest dry matter content on the list. High starch content. Late maturing variety. Bred by Limagrain Advanta Nederland BV, The Netherlands. Irl Agent: Seed Technology Ltd.
- Beacon (R):** **Uncovered:** Good yielding variety with good dry matter content. Very high starch content. Medium-early maturing variety. Bred by Limagrain Verneuil Holding, France, Irl Agent: Goldcrop Ltd.
- Beethoven (R):** **Uncovered:** High yielding variety. Moderate dry matter content. Low starch content. Medium-late maturing variety. Bred by Limagrain Verneuil Holding, France. Irl Agent: Seed Technology Ltd.
- Kroft (R):** **Uncovered:** Below average yielding variety. Very high Dry Matter and exceptional starch content, the highest on the Recommended List. A very early maturing variety and suitable for less favourable sites. Bred by KWS, Germany. Irl Agent: Seed Technology Ltd.
- Severus (PR-1):** **Uncovered:** First entered the list in 2015 with a provisional recommendation. Yield and dry matter are good and starch content is very high. An early maturing variety. Bred by KWS, Germany. Irl Agent: Seed Technology Ltd.

RECOMMENDED LISTS

- Cereal Varieties
- Herbages Varieties: (Grasses and Clover)
- Forage Maize Varieties
- Spring Bean Varieties
- Winter Oilseed Rape Varieties

CROPS SCHEMES AND SERVICES

- Seed Certification
- Seed Testing

The use of certified seed ensures a high level of varietal purity and germination.

Requests for this booklet should be sent to:

Crop Policy, Production & Safety Division,
Department of Agriculture, Food and the Marine,
Administrative Building, Backweston Campus,
Stacumny Lane, Celbridge, Co Kildare.

or

E-mail: annl.murphy@agriculture.gov.ie

Alternatively, Recommended Lists can be obtained from the Department of Agriculture, Food and the Marine website; www.agriculture.gov.ie where they can be found under the heading 'PUBLICATIONS'.



**Department of Agriculture, Food and the Marine
An Roinn Talmhaíochta, Bia agus Mara**

Agriculture House
Kildare Street
Dublin 2
Ireland

Áras Talmhaíochta
Sráid Chill Dara
Baile Átha Cliath
Éire

Telephone +353 (0)1 607 2000
Fax +353 (0)1 661 6263
info@agriculture.gov.ie

www.agriculture.gov.ie